References:-

[1] Abdel-Hafez, (2007), Benzimidazole condensed ring systems: New synthesis and antineoplastic activity of substituted 3,4-dihydro- and 1,2,3,4-tetrahydro- benzo [4,5]imidazo[1,2-a]pyrinnidine derivatives , *Archives of Pharmacal Research*., **30(6)**, 678-684.

[2] Abdel-Rahman, A.H., Keshk, E.M. Hanna, M.A., El-Bady, S.M., (2004), Synthesis and evaluation of some new spiro indoline-based heterocycles as potentially active antimicrobial agents. *Bioorg. Med. Chem.*, **12**, 2483–2488.

[3] Mobinikhaledi A., Foroughifar N. and Khodaei H., (2010), Synthesis of octahydroquinazoline derivatives using silica sulfuric acid as an efficient catalyst, *European Journal of Chemistry*, **1(4)**, 291-293.

[4] Chimirri A., Sarro A.D., Sarro G.D., Grasso S., Trimarchi G.R., and Zappall M., (1989), Synthesis and Anticonvulsant Properties of 2,3,3a,4-Tetrahydro- 1H-pyrrolo[1,2-a Ibenzimidazol- 1-ones, *J. Med. Chem*, **32**, 93-95.

[5] Stewart A.O., Bhatia P.A., McCarty C.M., Patel M.V., Staeger M.A., Arendsen D.L., Gunawardana I.W., Melcher L.M.,Dong Zhu G., Boyd S.A., Fry D.G., Cool B.L., Kifle L., Lartey K., Marsh K.C., Anita J., Kilgannon K.P., Wendy W. W., Meyer J. W., Gallatin M., and Okasinski G.F., (2001), Discovery of Inhibitors of Cell Adhesion Molecule Expression in Human Endothelial Cells. 1. Selective Inhibition of ICAM-1 and E-Selectin Expression, *J. Med. Chem***, 44**, 988-1002.

[6] Chawla A., Kaur R. and Goyal A., (2011), Importance of Microwave Reaction on the synthesis of Novel Benzimidazole Derivatives:A Review, *J. Chem. Pharm. Res*., **3(6)**, 925-944.

[7] B. E. Evans, K. E. Rittle, M. G. Bock, R. M. DiPardo, R. M. Freidinger, W. L. Whitter, G. F. Lundell D. F. Veber, P. S. Anderson, R. S. L. Chang, V. J. Lotti, D. J. Cerino, T. B. Chen, P. J. Kling, K. A. Kunkel J. P. Springer,and J. Hirshfieldt, (1988), Methods for Drug Discovery: Development of Potent, Selective, Orally Effective Cholecystokinin Antagonists, *J. Med. Chem*., **31**, 2235-2246.

[8] Bishop J.F, (1998), Approaches to induction therapy with adult acute myeloid leukaemia, *Acta Haematol*, **99**, 133-137.

[9] Boutaleb-Charki S., Marin C., Maldonado C.R., Rosales M.J., Urbano, J., Guitierrez- Sanchez R., Quiros M., Salas J.M., Sanchez-Moreno M., (2009),

Copper (II) complexes of [1,2,4]triazolo [1,5-a] pyrimidine derivatives as potential anti-parasitic agents. *Drug. Metab. Lett.*, **3**, 35–44.

[10] Piquani B. and Zhang W., (2011), Synthesis of diverse dihydropyrimidine-related scaffolds by fluorous benzaldehyde-based Biginelli reaction and post-condensation modifications, *Beilstein J. Org. Chem.*, **7**, 1294–1298.

[11] Cochrane Review, (2002), Multi-agent chemotherapy for early breast cancer, *Cochrane Database Syst Rev*, CD000487.

[12] Rich D.H., Moon B.J., and Harbeson S., (1984), Inhibition of Aminopeptidases by Amastatin and Bestatin Derivatives. Effect of Inhibitor Structure on Slow-Binding Processes, *J.m.chem.*, **27(4)**, 417-422.

[13] Patil D.R., Salunkhe S.M.,Deshmukh M.B. and Anbhule P.V., (2010), Synthesis of 6-Amino-5-cyano-4-Phenyl-2-Mercapto Pyrimidine using Phosphorous pentoxide, *The open Catalysis Journal*, **3** , 83-86.

[14] El-GendyM.M.A., Shaaban M., Shaaban K.A., El-Bondkly A.M., Laatsch H. Essramycin, (2008), A first triazolopyrimidine antibiotic isolated from nature. *J. Antibiot.*, **61**, 149–157.

[15] Akbas E., Levent A., Gumus S., Sumer M.R., and Akyazı I., (2010), Synthesis of Some Novel Pyrimidine Derivatives and Investigation of their Electrochemical Behavior, *Bull. Korean Chem. Soc.*, **31(12)**, 3622-3638.

[16] Fischer, G., (2008), Recent progress in 1,2,4-triazolo[1,5-a]pyrimidine chemistry. *Adv. Heterocycl. Chem.*, **95**, 143–219.

[17] Galmarini C.M, Mackey J.R, (1991) Mechanisms of drug resistance and reversal strategies. Leukemia, *Dumontet C. Nucleoside analogues,* **15**, 875-890.

[18] Harari P.M., (1997), Why has introduction chemotherapy of advanced head and neck cancer become a United States community standard of practice? *J Clin Oncol*, **15**, 2050-2055.

[19] Wahe.H., Asobo P.F., Cherkasov R.A., Wahe Z.H.T, Fomum Z.T. and Doepp D., (2004), Heterocycles of biological importance: Part-8 Formation of pyrimido[1,2-*a*]benzimidazoles and oxazolo[3,2-*a*] benzimidazoles by conjugate addition of 2-aminobenzimidazoles to 4-hydroxy-2-alkynenitriles , *Arkivoc,* **(1)**, 130-137.

[20] Roberge J.Y., yu G., Mikkillineni A., Wu X., Zhu Y., Lawrence R.M. and Ewing W.R., (2007), Synthesis of Triazolopyridines and triazolopyrimidines using a modified Mitsunobu reaction, *Arkivoc*, **(12)**,132-147.

[21] Jha M., Guy S., Ting Y.C., (2011), Microwave assisted synthesis of indole-annulated dihydropyrano[3,4-c]chromene derivatives via hetero-Diels–Alder reaction. *Tetrahedron Lett.*, **52**, 4337–4341.

[22] Kalisiak J.,Sharpless K.B. and Fokin V.V., (2008), Efficient Synthesis of 2-Substituted-1,2,3-triazoles, *Org. Lett.*, **10(15)**, 3171.

[23] Krasovsky, A. L., Moiseev, A. M., Nenajdenko, V. G., Balenkova, E. S., (2004), Synthesis of new fluorine containing triazolo- and tetrazolopyrimidines. *Synthesis-Stuttgart*, **7**, 901-905.

[24] Macdonalds J.S, Astrow A.B., (2001), Adjuvant therapy of colon cancer, *Semin Oncol*, **28**, 30-40.

[25] Pryadeina M.V., Burgart Y.V., Saloutin V.I., Chupakhin O.N. , (2008), Ring-chain isomerism of ethyl 7-polyfluoroalkyl-7-hydroxy-4,7-dihydro[1,2,4]triazolo[1,5-*a*]pyrimidine-6-carboxylates, *Mendeleev Comunications*, **18(5)**, 276-277.

[26] Martin M.W., Newcomb J., Nunes J.J, Boucher C., Chai L., Epstein L.F., Flores T.S., Gallant P., Gore A., Gu Y., Hsieh F., Huang X., Kim J.L., Morgenstern S.K., Antonio Oliveira-dos-Santos, Patel V.F, Powers D., Rose O., Tudor Y., Turci O.M., Welcher A.A., Zack D., Zhao H., Zhu L., Zhu X., Ghiron C., Monika Erman David Johnston, and Carl-Gustaf Pierre Saluste, (2008), Structure-Based Design of Novel 2-Amino-6-phenyl-pyrimido[5′,4′:5,6]pyrimido[1,2-*a*]benzimidazol 5(6*H*)-ones as Potent and Orally Active Inhibitors of Lymphocyte Specific Kinase (Lck) Synthesis SAR, and In Vivo Anti-Inflammatory Activity, *J. Med. Chem.*, **51**, 1637–1648.

[27] Buddh M.B., Bapodra A.H. and Ladva K.D., (2011), Synthesis and biological evaluation of thiazolo [3, 2-a] pyrimidine derivatives as a new type of potential antimicrobial agents. *Rasayan J. Chem*., **4(4)**, 824-828.

[28] Elwan N.M., Awad E.M., Hassaneen H.M., Anthony Linden, Heinz Heimgartner, (2003), Synthesis of 1,2,4-Triazolo[4,3-*a*]pyrimidine Derivatives by Cyclocondensation of a 2-Thioxopyrimidin-4(3*H*)-one with Hydrazonoyl Halides. *HCA*, **86**, 739–749.

[29] Pinedo H.M., Peters G.F.., (1988), Fluorouracil, biochemistry and pharmacology. *J Clin Oncol*, **6**, 1653-1664.

[30] Rai, U.S., Isloor A.M., Shetty P., Vijesh A.M., Prabhu N., Isloor S. Thiageeswaran M., Hoong-Kun F., (2010), Novel chromeno [2,3-b]pyrimidine derivatives as potential anti-microbial agents, *Eur. J. Med. Chem.*, **45**, 2695–2699.

[31] Kumbhre R.M., Kosurkar U., K. Vijay Kumar, Dadmal T., Tiwari A.K., (2010), Synthesis of multiplebiologically active 1,2-Dihydropyrimido[1,2-A]benzimidazole-3-carbonitrile, *IJABPT*, **1(2)**, 729-735.

[32] Ruisi G., Canfora L., Bruno G., Rotondo A., Mastropietro T.F., Debbia E.A., Girasolo M.A., Triorganotin M.B., (2010), Derivatives of 7-amino-2-(methylthio)-[1,2,4]triazolo[1,5-a] pyrimidine-6-carboxylic acid. Synthesis, spectroscopic characterization, in vitro antimicrobial activity and X-ray crystallography, *J. Organomet. Chem.*, **695**, 546–551.

[33] Sagar R., Moon-Ju K., Park S.B., (2008), An improved synthesis of pyrimidine- and pyrazole-based acyclo-C-nucleosides as carbohybrids, *Tetrahedron Lett.*, **49**, 5080–5083.

[34] Salameh S., Abul-Haj M., Quiros M., Salas J.M., (2005), 1,2,4-Triazolo[4,3-a]pyrimidines: A new kind of ligands. Structure of the silver (I) dimer with the 7-oxo derivative. *Inorg. Chim. Acta*, **358,** 824–827.

[35] Salgado A., Varela C., Collazo A.M.G., Pevarello P., (2010), Differentiation between[1,2,4]triazolo[1,5-a] pyrimidine and [1,2,4]triazolo[4,3-a]-pyrimidine regioisomers by 1H-15N HMBC experiments, *Magn. Reson. Chem.*, **48**, 614–622.

[36] El-moghazy S.M. , Diaa A. Ibrahim **,** Nagwa M. Abdelgawad, Nahla A. H. farag, (2011), Design , Synthesis and Biological Evaluation of Novel Pyrimido[4,5-d]pyrimidine CDK2 Inhibitors as Anti-Tumor Agents, *Sci. Pharm.*, **79**, 429–447.

[37] Salameh S., Abul-Haj M., Quiros M., Salas J.M., (2005), 1,2,4- Triazolo[4,3-a] pyrimidines: A new kind of ligands. Structure of the silver(I) dimer with the 7-oxo derivative, *Inorganica Chimica Acta*, **358**, 824–827.

[38] Shaaban M.R., Saleh T.S., Farag A.M., (2007), Synthesis and antimicrobial evaluation of novel pyrazolo[1,5-a] pyrimidine, triazolo[1,5-a]pyrimidine and pyrimido[1,2-a]benzimidazole derivatives, *Heterocycles*, **71**, 1765–1777.

[39] Sorenson S., Glimelius B., Nygren P., SBU group, (2001), Swedish council of Technology Assessment in Health Care. A systematic overview of chemotherapy effects in non-small cell lung cancer, *Acta Oncol*, **40**, 327-339.

[40] Zhang W., Williams J.P., Lu Y., Nagashima T. and Chu Q.**,** (2003), Fluorous synthesis of sclerotigenin-type benzodiazepine-quinazolinones *Chem. Rev*, **103**, 893.

[41] Wiernik P.H.., (2000), Current status of and futureprospects for the medical management of adenocarcinoma of the exocrine pancreas, *J Clin Gastroenterol*, **30**, 357-363.

[42] Wendelin W., Gobinitzer E. and Dalal abou El Ella, (2000), Synthesis and structure elucidation of Pyrimidobenzimidazoles and Fused derivatives II. Dodecahydrobenzimidazo [2,1-b]quinazolineand odecahydrobenzimidazo[2,1-b]benzo[h] quinazolines, *Monatshefte fur Chemie*, **131**, 353-374.

[43] William Engen a, Terrence E. O'Brien a, Brendan Kelly a, Jacinda Do a, Liezel Rillera a, Lance K. Stapleton Jack F. Youngren b, Marc O. Anderson (2010) Synthesis of aryl-heteroaryl ureas (AHUs) based on 4-aminoquinoline and their evaluation against the insulin-like growth factor receptor (IGF-1R), *Bioorganic & Medicinal Chemistry*, **18**, 5995–6005.

[44] Chang-Sheng Y., Song L., Cui-Hua W., Chen-Xia Y., Qing-Qing S., Shu-Jiang T., (2008), One- pot Three-component Solvent-free Synthesis of Benzo[4,5]imidazo[1,2-a]pyrimidine Derivatives Catalyzed by Sulfamic Acid, *Chinese Journal of Chemistry*, **26**, 2107-2111.